ForPatients

by Roche

Triple Negative Breast Cancer Breast Cancer

A Study of Cobimetinib Plus Paclitaxel, Cobimetinib Plus Atezolizumab Plus Paclitaxel, or Cobimetinib Plus Atezolizumab Plus Nab-Paclitaxel as Initial Treatment for Participants With Triple-Negative Breast Cancer That Has Spread

A Study of Cobimetinib Plus Paclitaxel, Cobimetinib Plus Atezolizumab Plus Paclitaxel, or Cobimetinib Plus Atezolizumab Plus Nab-Paclitaxel as Initial Treatment for Participants With Triple-Negative Breast Cancer That Has Spread

Trial Status Trial Runs In Trial Identifier
Terminated 13 Countries NCT02322814 2014-002230-32
WO29479

The source of the below information is the publicly available website ClinicalTrials.gov. It has been summarised and edited into simpler language.

Trial Summary:

This three-cohort, multi-stage, randomized, Phase II, multicenter trial will evaluate the safety and tolerability and estimate the efficacy of cobimetinib plus paclitaxel versus placebo plus paclitaxel in Cohort I, of cobimetinib plus atezolizumab plus paclitaxel in Cohort III in participants with metastatic or locally advanced, triple-negative adenocarcinoma of the breast who have not received prior systemic therapy for metastatic breast cancer (MBC). Participants may continue on study treatment until the development of progressive disease (PD) or the loss of clinical benefit, unacceptable toxicity, and/or consent withdrawal. The Cohort I target sample size is 12 participants for the safety run-in stage and approximately 90 participants in the expansion stage. Each of Cohorts II and III will consist of a safety run-in stage of approximately 15 participants.

Hoffmann-La Roche Sponsor		Phase 2 Phase	
NCT02322814 2014-002230-32 WO29479 Trial Identifiers			
Eligibility Criteria:			
Gender	 Age		Healthy Volunteers

ForPatients

by Roche

All >= 18 Years No